

On a conjecture about repdigits in k –generalized Fibonacci sequences

JHON J. BRAVO¹

UNIVERSITY OF CAUCA, COLOMBIA
MATHEMATICAL INSTITUTE, UNAM, MEXICO CITY

ARCTIC NUMBER THEORY WORKSHOP
SAARISELKÄ LAPLAND FINLAND, 15–20 JUNE, 2013

Abstract

For an integer $k \geq 2$, we consider the k –generalized Fibonacci sequence $(F_n^{(k)})_n$ which starts with $0, \dots, 0, 1$ (k terms) and each term afterwards is the sum of the k preceding terms. F. Luca [2] in 2000 and recently D. Marques [3] proved that 55 and 44 are the largest numbers with only one distinct digit (so called *repdigits*) in the sequences $(F_n^{(2)})_n$ and $(F_n^{(3)})_n$, respectively. Further, Marques conjectured that there are no repdigits having at least 2 digits in a k –generalized Fibonacci sequence for any $k > 3$. In this talk, we report about some arithmetic properties of $(F_n^{(k)})_n$ and confirm the conjecture raised by Marques. This is a joint work with Florian Luca.

Key words and phrases: Fibonacci numbers, lower bounds for nonzero linear forms in logarithms of algebraic numbers, continued fractions, repdigits.

References

- [1] J. J. Bravo and F. Luca, *On a conjecture about repdigits in k –generalized Fibonacci sequences*, to appear in Publ. Math. Debrecen.
- [2] F. Luca, *Fibonacci and Lucas numbers with only one distinct digit*, Port. Math. **57** (2) (2000), 243–254.
- [3] D. Marques, *On k –generalized Fibonacci numbers with only one distinct digit*, to appear in Util. Math.

¹jhonjaba@gmail.com